# **ONE-DAY**

# Colonoscopy Preparation with SUPREP



ENDOSCOPY CENTER	ENDOSCOPY (HOSPITAL)	APPOINTMENT DATE
17820 Talbot Rd S Renton, WA 98055	400 South 43rd St Renton, WA 98055	
425.690.3636	Park in South Tower Garage (Parking G). On Level P-2, enter the	CHECK IN TIME
Park in lot at the center.	Special Procedure Care Unit (SPCU) reception area and check in.	
<b>Medications and Precautions</b>	·	
<ul> <li>No anti-inflammatories one day before procedure such as Advil (ibuprofen) or Aleve (naproxen)—         Tylenol (acetaminophen) is okay.</li> <li>□ Low fiber diet, no multi-vitamins, iron, calcium supplements, seeds, or nuts for 7 days before procedure.</li> </ul>	Hold GLP-1 receptor agonists used for diabetes or weight loss, such as <b>semaglutide</b> (Ozempic, Rybelsus, Wegovy), <b>liraglutide</b> (Victoza, Saxenda), <b>dulaglutide</b>	IMPORTANT: BECAUSE THIS IS A SEDATED PROCEDURE, YOU MUST HAVE A DRIVER TAKE YOU HOME.  Buy Prescription Item at the Pharmacy
Continue heart and blood pressure medications and aspirin—unless your doctor tells you to stop.  Diabetes medicine—take 1/2 dose of insulin/oral diabetes medicine the evening before procedure. Hold insulin and/or oral diabetes medications the		SUPREP BOWEL PREP KIT   The state of control of the
day of the procedure.	of <b>24 hours</b> prior to procedure.	

# ONE DAY BEFORE PROCEDURE

1. DRINK ONLY CLEAR LIQUIDS THE ENTIRE DAY; NO SOLID FOODS; NO MILK PRODUCTS OR NON-DAIRY CREAMER; NO ALCOHOLIC BEVERAGES.

#### **APPROVED**

Beverages: tea, coffee, soda/carbonated beverages, fitness water, sports drinks

Fruit juices: apple, cran-apple, white grape,

lemonade (no pulp)

**Desserts:** flavored Jell-O, popsicles without pulp

Bouillon: clear, fat-free bouillon ONLY

Misc: salt, sugar





# **AVOID**

No RED or PURPLE colors

No solid food

**No** milk products or non-dairy creamer

No alcoholic beverages

No stock, broth, or soup (store bought or homemade)



## ONE DAY BEFORE PROCEDURE (continued)

- 2. As a comfort measure, apply Vaseline, A&D ointment, or Desitin to the anal area before laxative use.
- 3.

At about 6 PM on the evening before your procedure, complete Steps 4 through 7.





4. Pour ONE 6 oz. bottle of **SUPREP** liquid into the mixing container.



**5.** Add cool drinking water to the 16 oz. line on the container and mix.



6. Drink ALL the liquid in the container.



7. You MUST drink TWO more 16 oz. containers of water over the next one hour.

## **DAY OF PROCEDURE**

- \_\_\_ (6 hours before your scheduled procedure), repeat Steps 4 through 7 (above) using the other 6 oz. bottle of SUPREP.
- 2. It is normal to pass yellow/green liquid stool after the prep.
- 3. CLEAR LIQUIDS ONLY UP TO 2 HOURS BEFORE YOUR PROCEDURE CHECK-IN TIME, THEN NOTHING BY MOUTH.





WEralley VALLEY MEDICAL CENTER







Use the Google Translate app for your language