

AFTERNOON

Colonoscopy Preparation (Over-the-Counter)



UW Medicine
VALLEY
MEDICAL CENTER

☐ ENDOSCOPY CENTER

17820 Talbot Rd S
Renton, WA 98055

425.690.3636

Park in lot at the center.

☐ ENDOSCOPY (HOSPITAL)

400 South 43rd St
Renton, WA 98055

*Park in South Tower Garage
(Parking G). On Level P-2, enter the
Special Procedure Care Unit (SPCU)
reception area and check in.*

APPOINTMENT DATE

CHECK IN TIME

Medications and Precautions

☐ No anti-inflammatories one day before procedure such as Advil (ibuprofen) or Aleve (naproxen)—**Tylenol (acetaminophen) is okay.**

☐ Low fiber diet, no multi-vitamins, iron, calcium supplements, seeds, or nuts for **7 days before procedure.**

☐ Continue heart and blood pressure medications and aspirin—**unless your doctor tells you to stop.**

☐ **Diabetes medicine—take 1/2 dose of insulin/oral diabetes medicine the evening before procedure. Hold insulin and/or oral diabetes medications the day of the procedure.**

☐ Hold blood thinners such as _____ for _____ days before procedure.

☐ Hold GLP-1 receptor agonists used for diabetes or weight loss, such as **semaglutide** (Ozempic, Rybelsus, Wegovy), **liraglutide** (Victoza, Saxenda), **dulaglutide** (Trulicity), **tirzepatide** (Mounjaro), or _____ as follows:

- If you are on weekly dosing, your last dose must be a minimum of **one week** prior to procedure.
- If you are on daily dosing, your last dose must be a minimum of **24 hours** prior to procedure.

! IMPORTANT: BECAUSE THIS IS A SEDATED PROCEDURE, YOU MUST HAVE A DRIVER TAKE YOU HOME.

Buy These Items at the Pharmacy

☐ 8.3 oz. bottle of Miralax powder

☐ Dulcolax (bisacodyl) stimulant laxative tablets, 5 mg, 4 tablets total (NOT stool softeners)

☐ 64 oz. of Gatorade (one 64 oz. bottle OR two 32 oz. bottles) NO RED

☐ 10 oz. bottle of Magnesium Citrate (Lemon-Lime); **if you have congestive heart failure (CHF) or kidney/renal failure—do NOT buy**

THE DAY BEFORE YOUR PROCEDURE

1. YOU MAY EAT NORMALLY THROUGH LUNCH.

2. AFTER 4 PM, DRINK ONLY CLEAR LIQUIDS.



APPROVED

Beverages: tea, coffee, soda/carbonated beverages, fitness water, sports drinks

Fruit juices: apple, cran-apple, white grape, lemonade (no pulp)

Desserts: flavored Jell-O, popsicles without pulp

Bouillon: clear, fat-free bouillon **ONLY**

Misc: salt, sugar



AVOID

No RED or PURPLE colors

No solid food

No milk products or non-dairy creamer

No alcoholic beverages

No stock, broth, or soup (store bought or homemade)



3. As a comfort measure, apply Vaseline, A&D ointment or Desitin to the anal area before laxative use. It is normal to pass yellow/green liquid stool after the prep.



4. **At 7 PM**, mix together the **ENTIRE BOTTLE** (8.3 oz./ 238 g) of MiraLax with **64 oz.** of chilled Gatorade. You will drink this between 6 AM and 9 AM tomorrow.



5. **At 7 PM**, drink one 10 oz. bottle of Magnesium Citrate (Lemon-Lime). **Do NOT drink if you have congestive heart failure or kidney/renal failure.** You may mix it with Sprite, 7-up or ginger ale.




6. **At 9 PM**, take two 5 mg Dulcolax tablets with a glass of water.

DAY OF PROCEDURE



1. **At 5 AM**, take two 5 mg Dulcolax tablets with a glass of water.



2. **At 6 AM**, start drinking an 8 oz. glass of the MiraLax and Gatorade solution every 15 minutes until it is gone. Finish by 9 AM. It is normal to pass yellow/green liquid stool following the prep.

3. **CLEAR LIQUIDS ONLY UP TO 2 HOURS BEFORE YOUR PROCEDURE TIME, THEN NOTHING BY MOUTH.**



UW Medicine | VALLEY MEDICAL CENTER

